

Entrez PubMed Overview Help | FAQ Tutorials New/Noteworthy E-Utilities

PubMed Services
Journals Database
MeSH Database
Single Citation Matcher
Batch Citation Matcher
Clinical Queries
Special Queries
LinkOut
My NCBI

Related Resources
Order Documents
NLM Mobile
NLM Catalog
NLM Gateway
TOXNET
Consumer Health
Clinical Alerts
ClinicalTrials.gov
PubMed Central

A lymphotoxin-beta-specific receptor.

Science. 1994 Apr 29;264(5159):667-8.

Crowe PD, VanArsdale TL, Walter BN, Ware CF, Hession C, Ehrenfels B, Browning JL, Din WS, Goodwin RG, Smith CA.

Division of Biomedical Sciences, University of California, Riverside 92521.

Tumor necrosis factor (TNF) and lymphotoxin-alpha (LT-alpha) are members of a family of secreted and cell surface cytokines that participate in the regulation of immune and inflammatory responses. The cell surface form of LT-alpha is assembled during biosynthesis as a heteromeric complex with lymphotoxin-beta (LT-beta), a type II transmembrane protein that is another member of the TNF ligand family. Secreted LT-alpha is a homotrimer that binds to distinct TNF receptors of 60 and 80 kilodaltons; however, these receptors do not recognize the major cell surface LT-alpha-LT-beta complex. A receptor specific for human LT-beta was identified, which suggests that cell surface LT may have functions that are distinct from those of secreted LT-alpha.

Publication Types:

Comment on:

Comment

PMID: 8171323 [PubMed - indexed for MEDLINE]

Display /	Abstract	Show	20	Sort by	▼ Send	to 🔽

Write to the Help Desk

NCBI | NLM | NIH

Department of Health & Human Services

Privacy Statement | Freedom of Information Act | Disclaimer

Jan 19 2006 04:31:52